

Effects of g.10329C>T polymorphism in the SCD gene with milk production traits in Bangladeshi cattle: evidence from mutant protein function analysis

Md Abdul ALIM¹ (✉), Mridha Mohammad Kamal HOSSAIN¹, Nusrat JAHAN¹, Rubaya RUBAYA¹, Anjuman Ara BHUYAN¹, Jahangir ALAM¹, Ishtiaque AHAMMAD², Arittra BHATTACHARJEE², Mohammad Shohel Rana SIDDIKI³, Md MONIRUZZAMAN⁴

¹ Animal Biotechnology Division, National Institute of Biotechnology, Ganakbari, Ashulia, Savar, Dhaka-1349, Bangladesh

² Bioinformatics Division, National Institute of Biotechnology, Ganakbari, Ashulia, Savar, Dhaka-1349, Bangladesh

³ Department of Dairy Science, Bangladesh Agricultural University, Mymensingh -2202, Bangladesh

⁴ Molecular Biotechnology Division, National Institute of Biotechnology, Ganakbari, Ashulia, Savar, Dha-ka-1349, Bangladesh

✉ Corresponding author: alimmdcau@gmail.com

Received: May 11, 2024; Accepted: August 20, 2024

ABSTRACT

This study aimed to pinpoint the single nucleotide polymorphisms (SNPs) within the stearoyl-CoA desaturase (SCD) gene that is notably linked to milk production traits and to evaluate the structural and functional consequences of mutant proteins. Through pooled DNA sequencing, five SNPs were identified within the SCD gene. Among these, only one exonic SNP (g.10329C>T) was non-synonymous, resulting in an amino acid change from alanine to valine in the protein. Structural and functional analysis of the g.10329C>T mutation revealed discernible differences between wild-type and mutant proteins, indicating consequential effects on the observed phenotype. Subsequently, this identified SNP was genotyped in 100 milking cows for association analysis. The SNP g.10329C>T demonstrated significant allele substitution effects on milk yield traits. These findings suggest that the identified polymorphism (g.10329C>T) influences milk yield traits within the studied population and could serve as a genetic marker for cattle selection processes aimed at enhancing productivity with proper validation.

Keywords: Single nucleotide polymorphism, association, SCD, mutant proteins

INTRODUCTION

In Bangladesh, in recent decades a large number of local cattle have been replaced by crossbred cattle to improve milk production for higher demand. The cross-breeding has been started in 1936. The pure Hariana bulls were imported to upgrade the local cattle through natural services from India. In 1958 by Directorate of Livestock Services (DLS) started cross-breeding with the Artificial Insemination (AI) program which was strengthened in 1975-76 and to date it is going on. In the 1960s, Sindhi, Sahiwal and Tharparkar bulls were imported from Pakistan and in 1973, Friesian and Jersey bulls were imported from Australia to cross with indigenous cattle

for upgrading milk production. At present time, the cattle of Bangladesh are composed of pure local breeds and crossbreds (Hossain et al., 2002; Hamid et al., 2017). Local cattle in Bangladesh are small in size and horned, and they are used for ploughing all over the country. Even under ideal dietary, management, and environmental conditions, their production level is minimal (Hossain et al., 2002). The National Health Strategy recommends a daily milk ratio of 250 millilitres/person. The actual average daily consumption is 193 millilitres/person, less than the recommended intake (DLS, 2021). The low level of national production cannot meet the demand

of Bangladesh's growing population (DLS, 2021). In contrast to the world, the milk production growth rate in Bangladesh is very negligible due to a lack of proper selective breeding guidelines. Over the past five decades, Bangladesh has endeavoured to bridge the significant gap between milk production and demand (NLDP, 2007). The National Livestock Development Policy, under the Ministry of Fisheries and Livestock, has outlined a three-tiered approach to address this challenge: a) short-term objectives involve the insemination of the highest-performing crossbred Holstein-Friesian cows, which yield 10 kg or more of milk, under intensive management systems, utilizing imported semen from progeny-tested bulls; b) mid-term goals entail inseminating crossbred Holstein-Friesian cows that produce 6-10 kg of milk, reared within semi-intensive management systems, with semen sourced from 50% Holstein-Friesian bulls proven through progeny testing; c) long-term aspirations focus on inseminating native cows, managed under low-input production systems, with semen from superior progeny-tested or pedigreed bulls of local cattle breeds, aiming to enhance milk production. This strategic framework aims to gradually enhance milk production to meet the nation's growing demand, addressing both immediate and future needs within the dairy industry (NLDP, 2007).

Milk, rich in protein, lipids, lactose, and calcium, is a top dietary source of nutrition. It contains essential micronutrients and bioactive components like vitamins, minerals, immunoglobulins, and enzymes, which play an important role in the newborn's health and development (Medrano et al., 2000). Thus, milk and dairy products are vital for a balanced diet (Pereira, 2014). The fat content and composition of milk influence the quality of dairy products (Chilliard et al., 2003). Excess carbohydrates are converted into fatty acids through *de novo* lipogenesis in the liver, adipose tissue, and mammary glands, forming triglycerides stored as energy reserves in animals (Berg, 2019). According to published studies by Mele et al. (2007) and Moioli et al. (2007), the differences in mammary SCD activity, linked to either polymorphism of SCD and regulation of expression or differences in downstream factors that would affect interactions

between enzymes and substrates, may be the vital reason for diet-independent variations in milk fat's conjugated linoleic acid (CLA) content (Peterson et al., 2002). The production of monounsaturated fatty acids (MUFA) can occur through several pathways; a key pathway involves the process of fatty acid synthesis, which takes place in the cytoplasm of animal cells and is regulated by the SCD enzyme. The key step is the desaturation process, where SCD introduces a double bond into the fatty acid chain at a specific position (between carbons 9 and 10). This converts a saturated fatty acid into a MUFA (Shimakata and Mihara, 1972; Strittmatter et al., 1974; Enoch et al., 1976). Because of biosynthesis function, numerous independent studies have already examined the associations between polymorphisms in a number of genes, including stearoyl-CoA desaturase (SCD) and milk traits (Hayes and Goddard, 2001; Grisart et al., 2002; Khatib et al., 2007; Khatib et al., 2008; Ozkan and Yakan, 2020; Ahsani et al., 2022; Azis and Anggraeni, 2023). Several cattle breeds have reported associations between a missense polymorphism in exon 5 (GenBank AY241932: g.10329CAAla > TVal) and the concentration of certain individual fatty acid and fatty acid unsaturation indices in the milk fat (Mele et al., 2007; Moioli et al., 2007; Schennink et al., 2008; Alim et al., 2012; Houaga et al., 2018; Ma et al., 2021).

Additionally, research on the SCD gene in cattle has led to the discovery of 8 SNPs that together constitute 2 haplotypes (A and B) (Medrano et al., 2000). The animal SCD system consists of four hydrophobic transmembrane domains that are anchored in the endoplasmic reticulum membrane (Buist, 2004). This literature reported (Buist, 2004) that the SCD protein, a key enzyme in the production of fatty acids and a mitochondrial protein, has a transmembrane nature. Maryam et al. (2016) reported that p.K158I (Ala>Val) in SCD is localized close to the steroid-binding domain, a functionally active area involved in fatty acid synthesis. They also hypothesized that since this variant changes hydrophilicity to hydrophobicity, it may facilitate transport across the mitochondrial membrane and boost fatty acid production or that the presence of this substitution in the vicinity of the steroid-binding

domain may alter protein functionality. This hydrophilic to hydrophobic substitution may have contributed to the rise in milk fat content in cows (Maryam et al., 2016). The biosynthesis function and previous association effects on milk fat suggest that the *SCD* gene may also affect production traits like milk yield, fat percentage, and protein percentage, which are the main breeding goals in current selection schemes of dairy cattle. In this study, we chose the *SCD* gene as a potential candidate gene for milk production traits, considering both its biological role and genomic location. The objective was to investigate SNPs within the *SCD* gene, analyze the protein structure to understand the implications of these genetic variations and evaluate their correlation with milk production traits in Bangladeshi Local and Holstein Cross cows. This research aims to potentially utilize of identified SNPs as genetic markers in cattle selection processes.

MATERIALS AND METHODS

Experimental animals and phenotypic data collection

A total of one hundred crossbred (Bangladeshi Local X Holstein) F1 generation cows were selected from Central Cattle Breeding and Dairy Farm (CCBDF), Savar, Dhaka those are reared in the same environment and management condition. Milk samples were collected from each lactating cow once in every lactation (within 90-100 days) period. After collection, milk samples were immediately transferred to the Animal Biotechnology Division of the National Institute of Biotechnology (NIB) to generate phenotypic data like protein and fat percentage using an auto milk analyzer (Lactoscan, Milk Analyzer, Bulgaria). Milk yield data from each milking cow (305 days' milk yield) was collected from CCBDF. The complete workflow employed in this study is shown in Figure 1.

Screening of polymorphisms and genotyping

Blood samples were collected from a chosen subset of cows (n = 100) for DNA extraction, which was performed using the TIANamp Blood DNA Kit following the manufacturer's guidelines. Primers designed according to Alim et al. (2012) and synthesized by Invitrogen (Invitrogen




Figure 1. Workflow employed for the study

Life Technologies, China) were utilized. A DNA pool was created (50ng/μl/animal) and subjected to sequencing to identify potential SNPs. PCR amplification of the pooled samples was conducted using a programmable thermal cycler (Biometra GmbH, Germany) and Invitrogen's DreamTaq Green PCR Master Mix. The 25 μL reaction mixture contained 50 ng of genomic DNA, 1 μl specific SCD gene primers, 12.5μl of premix and 8.5 μl ddH₂O. The amplification process included 5 min at 94 °C for initial denaturing followed by 35 cycles at 94 °C for 30 s; annealing at 58 (°C) for 30 s, extension at 72 (°C) for 30 s and a final extension at 72 (°C) for 7 min. Gel electrophoresis of PCR products was carried out using 2% agarose gel and Azure c150 gel imaging workstations to confirm amplification. Subsequently, PCR products were sequenced at the Molecular Biotechnology Division, NIB, using the ABI3500 sequencer (Applied Biosystems, USA).

Sequence data analysis was performed using BioEdit Sequence Alignment Editor version 7.0.9.0 and ClustalW multiple sequence alignment programs to detect genetic polymorphisms. Finally, the identified polymorphism (g.10329C>T) was genotyped in all animals using PCR and sequencing techniques.

Structural and functional Impact Prediction of SCD mutant protein associated with milk traits

Non-synonymous polymorphisms in the SCD gene were analyzed for their impact on protein structure and function, affecting milk phenotype. HHpred, a web-based tool (<https://toolkit.tuebingen.mpg.de/tools/hhpred>), was used to detect remote protein homology and predict structures, while PROCHECK (<https://servicesnmbi.ucla.edu/PROCHECK/>) evaluated structural integrity. SAVES web server and the Ramachandran plot confirmed structure quality. MUpro (<https://www.ics.uci.edu/~baldig/mutation.html>) predicted the effect of SNPs on protein stability using Support Vector Machines and Neural Networks. DynaMut2 analyzed protein motion, flexibility, and stability impact (Rodrigues et al., 2021). Conservation profile analysis was done with the Consurf server (Ashkenazy et al., 2016), and domain names were determined using the Pfam server (Apweiler et al., 2001). Molecular dynamics simulations of wild-type and mutant SCD proteins were performed using GROMACS for 100 ns (Abraham et al., 2015).

Statistical analysis

Pedigrees of the genotyped animals were traced back three generations, resulting in a total of 300 animals included in the analysis. The kinship matrix was calculated using MATLAB version 7.11.0.584. POPGENE software, version 1.32, was utilized to assess allelic and genotypic frequencies for the loci and to perform the Hardy-Weinberg equilibrium test. SAS 9.1.0 software (SAS Institute Inc. USA) was employed to estimate the genotypic effects on milk production traits. The analysis used the mixed procedure with an animal model (Lynch and Walsh, 1997):

$$Y = \mu + hys + L + G + \alpha + e,$$

where:

- Y is the phenotypic value;
- μ is the overall mean;
- hys is a herd-year-season effect;
- L is the fixed effect of lactation;
- G is a fixed effect corresponding to the genotype of polymorphisms;
- α is a random polygenic component for pedigree relationships;
- e is a random residual.

Bonferroni correction, adjusting the significance threshold to account for multiple comparisons, reducing the likelihood of false positives, was performed for multiple t-tests. The least squares mean was utilized for multiple comparisons to estimate the effects of SCD polymorphic genotypes on milk production traits. This likely involved comparing the means of different genotypic groups. Falconer and Mackay's equation was employed to compute the additive (a), dominance (d), and allele substitution (α) effects. These effects offer insights into how alleles at a specific locus influence the phenotype. Falconer and Mackay's (1996) equation was employed to compute the additive (a), dominance (d), and allele substitution (α) effects i.e.:

$$a = (AA - BB) / 2$$

$$d = AB - (AA + BB) / 2$$

$$\alpha = a + d (q - p)$$

where AA and BB are homozygous, AB is heterozygous genotype, and p and q are the allele frequencies.

RESULTS

Screening of single nucleotide polymorphisms and genotypes

In this study, we identified five SNPs within the SCD gene (GenBank: AY241932) through sequence analysis of pooled DNA samples. Among these SNPs, two (g.6926A>G and g.8646A>G) were situated within introns 3 and 4, while the remaining three (g.10153A>G, g.10213T>C, and g.10329C>T) were located in exon 5 (Figure 2).




Figure 2. SCD gene structure and genotyping of individual polymorphisms by chromatograph differences in studied animals

Specifically, only the g.10329C>T SNP was anticipated to induce an amino acid substitution, resulting in a change from alanine to valine at position 293 within the SCD protein. Table 1 illustrates the genotypic and allelic frequencies, along with the results of the Hardy-Weinberg equilibrium test (χ^2). The chi-square test demonstrated that the genotypic frequencies of the g.10329C>T locus within the population were consistent with Hardy-Weinberg equilibrium ($P>0.05$) (Table 2), suggesting less selection pressure at this particular site.

Table 1. PCR primers used for sequencing the SCD gene		
Exons	Sequence	Size of PCR product
Exon 1	L- GTTGGCAACGAATAAAAGAGG	384 bp
	R- CGCGGTGATCTCAACTCTTC	
Exon 2	L- GGACCGGGTCTATGCCTATC	552 bp
	R- CCATCCAGCCTCTCAGGAC	
Exon 3	L- GTTCCCTGGGACTCCTAAGC	499 bp
	R- CCGGAACCTTAACCACAAGGA	
Exon 4	L- GGCAACTCCATGACTTCTCC	594 bp
	R- CATGACCGTCCTAGGTCAAC	
Exon 5	L- CCCATTCGCTCTTGTCTGT	400 bp
	R- CGTGGTCTTGCTGTGGACT	
Exon 6	L- GCCTCTGAGGGGATCTATTTG	543 bp
	R- AGGCAGAGTTGTTGGCTTTC	

Source: Alim et al. (2012)

Table 2. Genotypic and allelic frequencies and Hardy–Weinberg equilibrium χ^2 test of SCD genotypes

Polymorphisms	Genotypic frequency			Allelic frequency		Hardy–Weinberg equilibrium χ^2 test
	CC	CT	TT	C	T	
g.10329C>T	0.23	0.65	0.12	0.55	0.45	$P > 0.05$

Structural and functional Impact Prediction of SCD mutant protein associated with milk traits

The three-dimensional model of the protein was generated (Figure 3) in PDB format using the HHPred server. The quality of the model was validated by inspecting the Ramachandran plot generated by the

PROCHECK web server. The findings indicated that the majority of the residues of the three-dimensional models belonged to the most favored regions for both the wild and mutant versions (Figure 4).




Figure 3. Three-dimensional model of SCD mutant g.10329 C>T (V293A) (the position of the mutated amino acid is marked in red)




Figure 4. Ramachandran plot of SCD mutant g.10329 C>T (V293A) generated by PROCHECK

The conservation profile findings were displayed in the form of a structural representation of the protein sequence, with the putative structural and functional

residues highlighted. Here, the V293A position in SCD belonged to the average conservation profile (Figure 5).




Figure 5. Conservation profile analysis of SCD protein

SCD contained only one domain namely Acyl-CoA desaturase and the SCD mutant V293A belonged to that domain (Figure 6). The Acyl-CoA desaturase domain is an essential component of desaturase enzymes, playing a

fundamental role in lipid homeostasis and cell function. This domain catalyzes the introduction of double bonds into fatty acyl chains of lipids, leading to the formation of unsaturated fatty acids.




Figure 6. SCD domains

Interatomic interactions prediction with Mupro analysis revealed that the mutation V293A in SCD decreased the stability of the protein. The delta G for mutations in V293A in SCD was -1.7005351 kcal/mol. Dynamut2 predicted stability change ($\Delta\Delta G_{\text{Stability}}$) for mutations in SCD was -1.74 kcal/mol. A negative value indicates a destabilizing effect of protein. The mutation was more unstable for the protein if the $\Delta\Delta G$ value was negative (Figure 7).

We calculated the RMSD to check the reliability of the systems. Variation in the RMSD value indicates a change in the protein. Throughout the simulation, the RMSD value between wild-type SCD and the mutant changed (Figure 8).

The regional flexibility of the protein was evaluated using the RMSF method which showed that the mutant protein was more flexible (Figure 9).




Figure 7. Interatomic interactions of SCD wild type and V293A mutant revealed by dyanmute2 analysis




Figure 8. RMSD of the wild type (green) SCD and the V293A mutant (red) (X-axis represents the time (ns) while the Y-axis represents the RMSD value (nm))




Figure 9. RMSF of the wild-type (green) SCD and the V293A (red) mutant (X-axis represents the amino acid residues while the Y-axis represents the RMSF value (nm))

The degree of compactness is measured using the radius of gyration. After 20 ns, the mutant protein had a higher radius of gyration compared to the wild type (Figure 10).




Figure 10. The radius of gyration of the wild-type SCD (green) and the V293A mutant (red) (the X-axis represents the time (ps) while the Y-axis represents the area (nm²))

The stability of proteins' hydrophobic cores was predicted using solvent-accessible surface area (SASA) in molecular dynamic simulations. Here, the SASA values declined gradually for both the wild type and the mutant. However, for the mutant, the SASA values remained higher than the wild type throughout the simulation period indicating a higher probability of disruption by solvents (Figure 11).




Figure 11. SASA of the wild-type SCD (green) and the V293A (red) mutant (X-axis represents the time (ps) while the Y-axis represents the SASA value (nm))

Association and effects of SNPs

The SNP (g.10329C>T) was initially identified as associated ($P \leq 0.0001$) with milk traits, with raw P values (<0.05). However, subsequent Bonferroni correction for multiple t-tests did not render these associations significant. Nevertheless, significant additive and allele substitution effects were observed for milk yield traits at the g.10329C>T loci ($P < 0.01$) (Table 3). The T allele demonstrated the potential to increase milk production by 198.46 kg over a full lactation period compared to the C allele at the g.10329C>T position (Table 3).

Table 3. Effects of g.10329 C>T polymorphism of different SCD genotypes on Bangladeshi local cattle (LSM $\pm$ SE)

Locus	Genotype (n)	Milk yield (kg)	Fat (%)	Protein (%)	Additive (a), Dominant (d) and Allele substitution (a) effects		
					Milk yield (kg)	Fat (%)	Protein (%)
g.10329 C>T	CC (23)	1946.37 $\pm$ 155.919	3.79 $\pm$ 0.097	3.07 $\pm$ 0.058	188.931 (a)*	0.029 (a)	-0.023 (a)
	CT (65)	1622.13 $\pm$ 90.020	3.59 $\pm$ 0.056	3.11 $\pm$ 0.027	-95.309 (d)	-0.078 (d)	0.018(d)
	TT (12)	1568.51 $\pm$ 209.188	3.64 $\pm$ 0.013	3.12 $\pm$ 0.020	198.462 (a)*	0.037 (a)	-0.025 (a)
P-value		<0.0001 (0.36-0.46 corrected)	<0.0001 (0.90 corrected)	<0.0001 (0.80 corrected)			
Maximum yield		3003.781	4.592	3.320		-	
Minimum yield		939.400	2.867	2.895		-	

Notes: * Significant effect at $P < 0.05$ level; n = number of animals

DISCUSSION

Stearoyl-coenzyme A desaturase (SCD), a multifunctional complex enzyme for cellular biosynthesis of MUFAs by catalyzing the introduction of a double bond into saturated fatty acyl-CoAs, has been studied for its association with milk production traits and milk fat-related traits in dairy cattle (Taniguchi et al., 2004; Conte et al., 2006; Mele et al., 2007; Macciotta et al., 2008; Schennink et al., 2008; Kgwatalala et al., 2009; Alim et al., 2012; Houaga et al., 2018; Ma et al., 2021). The primary mechanisms involved in milk fat synthesis include de novo synthesis of short-chain fatty acids (FASN, ACACA, and SCD), the uptake, transport, and activation of long-chain fatty acids (LPL, VLDLR, ACSL, FABP3, and SCD), triglyceride synthesis (GPAM, LPIN1, DGAT, and AGPAT6), signalling pathways, and the production of transcription factors (mTOR, SREBP, PPARG, and AMPK) (Guo et al., 2024). Macciotta et al.(2008) reported that Italian Holsteins cows with CC homozygous genotype at the SCD locus (g.10329C/T) produced higher milk and protein compared to homozygous TT cows; heterozygous CT cows were in an intermediate position. Similar to Macciotta et al.(2008), Taniguchi et al.(2004) also reported that cows with (A293V) AA genotypes producing higher milk fat compared to VV genotypes in Japanese Black cows. Interestingly Alim et al. (2012)

reported the association of heterozygous genotypes with different milk production traits in Chinese Holsteins and heterozygous CT cows produced more milk than the cows of homozygous CC or TT genotypes. Similar to Alim et al. (2012), Maryam et. al. (2016) reported that homozygous genotypes produced less fat than heterozygous genotypes in buffalo for the same SNP. Recently, it has been reported that in Polish Holstein-Friesian cows, the A293V (c.878C/T) mutation in SCD has been associated with milk fatty acids and altered the amino acid alanine to valine (Kesek-Wozniak et al., 2020). Constantly, Bouwman et al. (2011) documented that the A allele of SNP (A293V; c.878C/T) in SCD was linked to greater milk fatty acid levels, while other research indicated that the V allele had a less significant impact on milk fats in the White Fulani and Borgou cow breeds (Duchemin et al., 2013; Houaga et al., 2018). Very recent reports demonstrated that the SCD could be one of the candidates and g.10153A>G SNP (AG genotype) as a genetic marker to assist selection in reducing saturated fatty acid and increasing MUFA in the milk content of Holstein Friesian cows in Indonesia (Azis and Anggraeni, 2023). Our findings showed significant substitution effects and association at g.10329 C>T locus for milk yield trait. At g.10329 C>T locus, homozygous CC genotypes produced more milk compared to homozygous

TT genotypes and heterozygous AG genotypes were in an intermediate position but the result is inconsistent with Bayraktar and Ozcan (2023), they reported that heterozygous CT genotypes significantly associated with total milk yield similar to Alim et al. (2012).

The results of the mutational impact analysis conducted by Mupro suggest that mutations in *SCD* are associated with a destabilizing effect on the protein structure. This implies that these mutations could potentially disrupt the functionality of the protein to varying extents. The analysis of interatomic interactions further supports this idea, revealing discrepancies between the mutants and the wild type in terms of their interactions with neighbouring residues. Such differences indicate that the function of the proteins may be affected by these mutations. A protein's activity, regulation, and function rely heavily on the stability of its structure, any reduction in stability can lead to protein degradation, misfolding, and eventual dysfunction. This highlights the importance of comprehending the structural consequences of mutations in proteins like *SCD* (Rozario et al., 2021). Sequence conservation analysis unveiled that the mutation is located in average conserved locations suggesting that they might have important impacts on the functioning of the protein. The analysis of sequence conservation indicates that the mutation sites are situated in moderately conserved regions. This suggests that these mutations may exert significant effects on the protein's functioning. Conserved regions are typically crucial for protein structure and function, making mutations occurring in these regions more likely to have functional repercussions. Our result showed that *SCD* contained only the Acyl-CoA desaturase domain and the *SCD* mutant V293A was in the domain. The Acyl-CoA desaturase domain, also known as the fatty acid desaturase domain, is a conserved structural motif found in enzymes known as desaturases. These enzymes play a crucial role in lipid metabolism by introducing double bonds into fatty acyl chains. These double bonds are essential for the formation of unsaturated fatty acids, which are important components of various cellular structures, energy storage molecules, and signalling molecules. A molecular dynamics simulation complex lasting 100 nanoseconds

highlighted distinct variances between the wild-type *SCD* and the V293A mutant in terms of RMSD, RMSF, Radius of Gyration, and SASA profiles. This indicates that the mutation in the *SCD* protein is likely to disrupt its normal function. Consequently, it is plausible to predict that the mutation could affect both the structure and function of the protein, potentially influencing the observed phenotype. However, additional in vitro and in vivo experiments involving a larger number of animals and a more extensive dataset are imperative to validate our findings further.

CONCLUSIONS

The present research has unveiled significant effects of allele substitution related to milk yield traits at the g.10329C>T locus. Throughout the entire lactation period, the presence of the C allele correlated with a noteworthy increase of 198.46 kg in milk production compared to the T allele at the g.10329C>T location. Furthermore, we did a series of structural and functional analyses like structural prediction, interatomic interactions, conservation profile analysis, domain identification, and molecular dynamic simulation of mutant protein to find out the allele substitution effects. There have not been any previous reports with structural and functional analysis of mutant protein to support these findings. The association results along with structural and functional evaluation of the *SCD* mutant and wild protein, and useful information provided by this study could be used with proper validation in selective breeding initiatives aimed at enhancing the milk production performance of dairy cattle.

ACKNOWLEDGEMENT

The study was funded by the Bangladesh Academy of Sciences under BAS-USDA program Code No. LS-15/2017.

The authors would like to acknowledge the Department of Livestock Services, Bangladesh and the Bangladesh Academy of Sciences under the BAS-USDA program (Code No. LS-15/2017) for supporting the research. The authors also would like to thank a post-doctoral researcher at Wuhan University for his assistance with English expression and editing.

REFERENCES

- Abraham, M.J., Murtola, T., Schulz, R., Pall, S.R., Smith, J.C., Hess, B., Lindahl, E. (2015) GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX*, 1, 19-25.
DOI: <https://doi.org/10.1016/j.softx.2015.06.001>
- Ahsani, M.R., Mohammadabadi, M., Buchkovska, V., Levstafieva, Y.M., Kucher, D., Kochuk-Yashchenko, O., Babenko, O., Stavetska, R., Oleshko, V., Kalashnyk, O. (2022) Association of Stearoyl CoA Desaturase Expression with Cattle Milk Characteristics. *Iranian Journal of Applied Animal Science*, 12, 271-279.
- Alim, M.A., Fan, Y.P., Wu, X.P., Xie, Y., Zhang, Y., Zhang, S.L., Sun, D.X., Zhang, Q., Liu, L. (2012) Genetic effects of stearyl-coenzyme A desaturase (SCD) polymorphism on milk production traits in the Chinese dairy population. *Molecular Biology Reports*, 39, 8733-40.
DOI: <https://doi.org/10.1007/s11033-012-1733-6>
- Apweiler, R., Attwood, T.K., Bairoch, A., Bateman, A., Birney, E., Biswas, M., Bucher, P., Cerutti, L., Corpet, F., Croning, M.D.R. (2001) The InterPro database, an integrated documentation resource for protein families, domains and functional sites. *Nucleic acids research*, 29, 37-40. DOI: <https://doi.org/10.1093/nar/29.1.37>
- Ashkenazy, H., Abadi, S., Martz, E., Chay, O., Mayrose, I., Pupko, T., Ben-Tal, N. (2016) ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation in macromolecules. *Nucleic acids research*, 44, W344-W50.
DOI: <https://doi.org/10.1093/nar/gkw408>
- Azis, R., Anggraeni, A. (2023) Association of Stearoyl-CoA Desaturase Gene Polymorphisms on Milk Fatty Acid Composition of Holstein Friesian Cows in Indonesia. *Tropical Animal Science Journal*, 46, 27-35. DOI: <https://doi.org/10.5398/tasj.2023.46.1.27>
- Berg, J.S.L., Tymoczko, J., Gatto, G. (2019) *Biochemistry*. 9th ed. New York (NY): W.H. Freeman and Company.
- Bayraktar, M., Ozcan, B.D. (2023) Association between Stearoyl CoA Desaturase (SCD) Gene Polymorphisms and Milk Production in Holstein Cattle Breed. *Turkish Journal of Agriculture-Food Science and Technology*, 11, 799-802.
DOI: <https://doi.org/10.24925/turjaf.v11i4.799-802.5969>
- Bouwman, A.C., Bovenhuis, H., Visker, M.H.P.W., van Arendonk, J.A.M. (2011) Genome-wide association of milk fatty acids in Dutch dairy cattle. *BMC genetics*, 12, 1-12.
DOI: <https://doi.org/10.1186/1471-2156-12-43>
- Buist, P.H. (2004) Fatty acid desaturases: selecting the dehydrogenation channel. *Natural product reports*, 21, 249-62.
DOI: <https://doi.org/10.1039/B302094K>
- Chilliard, Y., Ferlay, A., Rouel, J., Lamberet, G. (2003) A review of nutritional and physiological factors affecting goat milk lipid synthesis and lipolysis. *Journal of Dairy Science*, 86, 1751-70.
DOI: [https://doi.org/10.3168/jds.S0022-0302\(03\)73761-8](https://doi.org/10.3168/jds.S0022-0302(03)73761-8)
- Conte, G., Mele, M., Castiglioni, B., Serra, A., Viva, M., Chessa, S., Pagnacco, G., Secchiari, P. (2006) Relationship between bovine SCD polymorphism locus and mammary gland desaturation activity. In: 8th World Congress on Genetics Applied to Livestock Production, pp. 22-37. Instituto Prociencia, Belo Horizonte, Minas Gerais, Brazil.
- DLS (2021) Available at: <https://www.dls.gov.bd/site/page/22b1143b-9323-44f8-bfd8-647087828c9b/Livestock-Economy>
- Duchemin, S., Bovenhuis, H., Stoop W.M., Bouwman, A.C., Van Arendonk, J.A.M., Visker, M. (2013) Genetic correlation between composition of bovine milk fat in winter and summer, and DGAT1 and SCD1 by season interactions. *Journal of Dairy Science*, 96, 592-604. DOI: <https://doi.org/10.3168/jds.2012-5454>
- Enoch, H.G., Catala, A., Strittmatter, P. (1976) Mechanism of rat liver microsomal stearyl-CoA desaturase. Studies of the substrate specificity, enzyme-substrate interactions, and the function of lipid. *Journal of Biological Chemistry*, 251, 5095.
DOI: [https://doi.org/10.1016/S0021-9258\(17\)33223-4](https://doi.org/10.1016/S0021-9258(17)33223-4)
- Falconer, D.S., Mackay, T.F.C. (1996) *Introduction to Quantitative Genetics*. 4th edn. Longman Scientific and Technical, New York.
- Grisart, B., Coppieters, W., Farnir, F., Karim, L., Ford, C., Berzi, P., Cambisano, N., Mni, M., Reid, S., Simon, P. (2002) Positional candidate cloning of a QTL in dairy cattle: identification of a missense mutation in the bovine DGAT1 gene with major effect on milk yield and composition. *Genome Research*, 12, 222-31.
DOI: <http://www.genome.org/cgi/doi/10.1101/gr.224202>
- Guo, Y., Wei, Z., Zhang, Y., Cao, J. (2024) Research Progress on the Mechanism of Milk Fat Synthesis in Cows and the Effect of Conjugated Linoleic Acid on Milk Fat Metabolism and Its Underlying Mechanism: A Review. *Animals*, 14, 204.
DOI: <https://doi.org/10.3390/ani14020204>
- Hamid, M., Rahman, A., Zaman, M.A., Hossain, K.M. (2017) Cattle genetic resources and their conservation in Bangladesh. *Asian Journal of Animal Sciences*, 11 (2), 54-64.
- Hayes, B., Goddard, M.E. (2001) The distribution of the effects of genes affecting quantitative traits in livestock. *Genetics Selection Evolution*, 33, 209-30.
DOI: <https://doi.org/10.1186/1297-9686-33-3-209>
- Hossain, K.B., Takayanagi, S., Miyake, T., Moriya, K., Bhuiyan, A., Sasaki, Y. (2002) Statistical genetic studies on cattle breeding for dairy productivity in Bangladesh: I. Genetic improvement for milk performance of local cattle populations. *Asian-Australasian Journal of Animal Sciences*, 15, 627-32.
DOI: <https://doi.org/10.5713/ajas.2002.627>
- Houaga, I., Muigai, A.W.T., Nganga, F.M., Ibeagha-Awemu, E.M., Kyallo, M., Youssao, I.A.K., Stomeo, F. (2018) Milk fatty acid variability and association with polymorphisms in SCD1 and DGAT1 genes in White Fulani and Borgou cattle breeds. *Molecular Biology Reports*, 45, 1849-62. DOI: <https://doi.org/10.1007/s11033-018-4331-4>
- Kesek-Wozniak, M.M., Wojtas, E., Zielak-Steciwo, A.E. (2020) Impact of SNPs in ACACA, SCD1, and DGAT1 genes on fatty acid profile in bovine milk with regard to lactation phases. *Animals*, 10, 997.
DOI: <https://doi.org/10.3390/ani10060997>
- Kgwatalala, P.M., Ibeagha-Awemu, E.M., Mustafa, A.F., Zhao, X. (2009) Influence of stearyl-coenzyme A desaturase 1 genotype and stage of lactation on fatty acid composition of Canadian Jersey cows. *Journal of Dairy Science*, 92, 1220-8.
DOI: <https://doi.org/10.3168/jds.2008-1471>
- Khatib, H., Monson, R.L., Schutzkus, V., Kohl, D.M., Rosa, G.J.M., Rutledge, J.J. (2008) Mutations in the STAT5A gene are associated with embryonic survival and milk composition in cattle. *Journal of Dairy Science*, 91, 784-93.
DOI: <https://doi.org/10.3168/jds.2007-0669>
- Khatib, H., Zaitoun, I., Wiebelhaus-Finger, J., Chang, Y.M., Rosa, G.J.M. (2007) The association of bovine PPARGC1A and OPN genes with milk composition in two independent Holstein cattle populations. *Journal of Dairy Science*, 90, 2966-70.
DOI: <https://doi.org/10.3168/jds.2006-812>
- Lynch, M., Walsh, B. (1997) *Genetics and Analysis of Quantitative Traits*. Sinauer Associates, Inc., Sunderland, MA, USA.
- Ma, Y., Khan, M.Z., Xiao, J., Alugongo, G.M., Chen, X., Chen, T., Liu, S., He, Z., Wang, J., Shah, M.K. (2021) Genetic markers associated with milk production traits in dairy cattle. *Agriculture*, 11, 1018.
DOI: <https://doi.org/10.3390/agriculture11101018>

- Macciotta, N.P., Mele, M., Conte, G., Serra, A., Cassandro, M., Dal Zotto, R., Borlino, A.C., Pagnacco, G., Secchiari, P. (2008) Association between a polymorphism at the stearoyl CoA desaturase locus and milk production traits in Italian Holsteins. *Journal of Dairy Science*, 91, 3184-9. DOI: <https://doi.org/10.3168/jds.2007-0947>
- Maryam, J., Babar, M.E., Bao, Z., Nadeem, A. (2016) A novel selection signature in stearoyl-coenzyme A desaturase (SCD) gene for enhanced milk fat content in *Bubalus bubalis*. *Tropical Animal Health and Production*, 48, 1343-9. DOI: <https://doi.org/10.1007/s11250-016-1092-8>
- Medrano, J.F., Johnson, A., DePeters, E.J., Islas, A. (2000) Genetic modification of the composition of milk fat: Identification of polymorphisms within the bovine stearoyl-CoA-desaturase gene. *Pflügers Archiv*, 439, R24-R.
- Mele, M., Conte, G., Castiglioni, B., Chessa, S., Macciotta, N.P., Serra, A., Buccioni, A., Pagnacco, G., Secchiari, P. (2007) Stearoyl-coenzyme A desaturase gene polymorphism and milk fatty acid composition in Italian Holsteins. *Journal of Dairy Science*, 90, 4458-65. DOI: <https://doi.org/10.3168/jds.2006-617>
- Moioli, B., Contarini, G., Avalli, A., Catillo, G., Orru, L., De Matteis, G., Masoero, G., Napolitano, F. (2007) Short communication: Effect of stearoyl-coenzyme A desaturase polymorphism on fatty acid composition of milk. *Journal of Dairy Science*, 90, 3553-8. DOI: <https://doi.org/10.3168/jds.2006-855>
- NLDP. (2007) National Livestock Development Policy, Ministry of Fisheries and Livestock, Government of the People's Republic of Bangladesh.
- Ozkan, H., Yakan, A. (2020) The relationship between milk fatty acid profile and expression levels of SCD, FASN and SREBP1 genes in damascus dairy goats. *Kocatepe Veterinary Journal*, 13, 294-303. DOI: <https://doi.org/10.30607/kvj.728554>
- Pereira, P.C. (2014) Milk nutritional composition and its role in human health. *Nutrition*, 30, 619-27. DOI: <https://doi.org/10.1016/j.nut.2013.10.011>
- Peterson, D.G., Kelsey, J.A., Bauman, D.E. (2002) Analysis of variation in cis-9, trans-11 conjugated linoleic acid (CLA) in milk fat of dairy cows. *Journal of Dairy Science*, 85, 2164-72. DOI: [https://doi.org/10.3168/jds.S0022-0302\(02\)74295-1](https://doi.org/10.3168/jds.S0022-0302(02)74295-1)
- Rodrigues, C.H.M., Pires, D.E.V., Ascher, D.B. (2021) DynaMut2: Assessing changes in stability and flexibility upon single and multiple point missense mutations. *Protein Science*, 30, 60-9. DOI: <https://doi.org/10.1002/pro.3942>
- Rozario, L.T., Sharker, T., Nila, T.A. (2021) In silico analysis of deleterious SNPs of human MTUS1 gene and their impacts on subsequent protein structure and function. *PloS one*, 16, e0252932. DOI: <https://doi.org/10.1371/journal.pone.0252932>
- Schennink, A., Heck, J.M., Bovenhuis, H., Visker, M.H., van Valenberg, H.J., van Arendonk, J.A. (2008) Milk fatty acid unsaturation: genetic parameters and effects of stearoyl-CoA desaturase (SCD1) and acyl CoA: diacylglycerol acyltransferase 1 (DGAT1). *Journal of Dairy Science*, 91, 2135-43. DOI: <https://doi.org/10.3168/jds.2007-0825>
- Shimakata, T., Mihara, K. (1972) Reconstitution of hepatic microsomal stearoyl-coenzyme A desaturase system from solubilized components. *Journal of Biochemistry*, 72, 1163-74. DOI: <https://doi.org/10.1093/oxfordjournals.jbchem.a130004>
- Strittmatter, P., Spatz, L., Corcoran, D., Rogers, M.J., Setlow, B., Redline, R. (1974) Purification and properties of rat liver microsomal stearyl coenzyme A desaturase. *Proceedings of the National Academy of Sciences*, 71, 4565. DOI: <https://doi.org/10.1073/pnas.71.11.456>
- Taniguchi, M., Utsugi, T., Oyama, K., Mannen, H., Kobayashi, M., Tanabe, Y., Ogino, A., Tsuji, S. (2004) Genotype of stearoyl-coA desaturase is associated with fatty acid composition in Japanese Black cattle. *Mammalian Genome*, 15, 142-8. DOI: <https://doi.org/10.1007/s00335-003-2286-8>

Original Article

Effects of *FASN* gene polymorphism on milk production traits in Bangladeshi cattle: Insights from mutant protein function

Alim, M. A.^{1*}; Hossain, M. M. K.¹; Jahan, N.^{1,2}; Rubaya, R.¹; Bhuyan, A. A.¹; Alam, J.¹; Ahammad, I.³; Siddiki, M. S. R.⁴ and Moniruzzaman, M.⁵

¹Animal Biotechnology Division, National Institute of Biotechnology, Ganakbari, Ashulia, Savar, Dhaka-1349, Bangladesh;

²Department of Animal Nutrition, Genetics and Breeding, Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka 1207, Bangladesh; ³Bioinformatics Division, National Institute of Biotechnology, Ganakbari, Ashulia, Savar, Dhaka-1349, Bangladesh;

⁴Department of Dairy Science, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh; ⁵Molecular Biotechnology Division, National Institute of Biotechnology, Ganakbari, Ashulia, Savar, Dhaka-1349, Bangladesh

*Correspondence: M. A. Alim, Animal Biotechnology Division, National Institute of Biotechnology, Ganakbari, Ashulia, Savar, Dhaka-1349. E-mail: alimmdcau@gmail.com



10.22099/ijvr.2025.49855.7351

(Received 31 Mar 2024; revised version 12 Jan 2025; accepted 2 Feb 2025)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Abstract

Background: To meet up the demand gap of milk in Bangladesh, short-term, midterm, and long-term goal have been set up by the government through crossing with Bangladeshi local cattle and high-producing foreign cattle like Friesian, Jersey, Sahiwal, etc. **Aims:** The purpose of this study was to identify the single nucleotide polymorphisms (SNPs) in the *FASN* gene, to check the structural and functional impact of mutant proteins on milk production traits that are significantly associated. **Methods:** Four SNPs were identified in exons 26, 36, 38, and 41 of the *FASN* gene using pooled DNA sequencing, but only one SNP g.17924 A>G was a non-synonymous that changed the amino acid threonine to alanine in the FASN protein and the other three SNPs were silent mutations. Structural and functional prediction analysis were done with a series of techniques to detect remote protein homology and predict structures, structural integrity, structure quality, protein stability, protein motion, flexibility, and stability impact, conservation profile and finally molecular dynamics simulations for wild-type and mutant protein expression differences. **Results:** The non-synonymous g.17924 A>G mutation showed a clear difference between wild and mutant proteins, indicating the impact on the observed phenotype. Then, SNP g.17924 A>G was genotyped in 100 milking cows aiming to check the association effects. SNP g.17924 A>G was found to have significant allele substitution effects on milk yield traits. **Conclusion:** Our results suggest that the identified polymorphism affects milk yield traits in the studied population and could be used as genetic marker for cattle selection processes aiming to increase productivity.

Key words: Association, Bangladeshi cattle, Candidate gene, *FASN*, Single nucleotide polymorphism

Introduction

Crossbred cattle have largely replaced native cattle in Bangladesh over recent decades to meet higher milk demand. Crossbreeding began in 1936 with imported Haryana bulls from India to upgrade the local cattle. In 1958, the Directorate of Livestock Services (DLS) launched artificial insemination (AI), a program that was expanded in 1975-76 and is still in operation today. In the 1960s, bulls from Pakistan (Sindhi, Tharparkar, Sahiwal) were imported, followed by Friesian and Jersey bulls from Australia in 1973, to enhance milk production. Bangladesh's cattle now consist of both crossbreeds and pure local breeds (Hossain *et al.*, 2002; Hamid *et al.*, 2017). Native cattle are small, horned, and used mainly for plowing, but their production is low

even under ideal conditions (Hossain *et al.*, 2002). The recommended daily milk intake is 250 ml/person, but actual consumption averages only 193 ml/person, failing to meet national recommended demand (DLS, 2021). For over 50 years, Bangladesh has struggled to close this gap because of the lack of effective selective breeding guidelines. Short-term (inseminate the top most cross bred Holstein-Friesian cows producing 10 kg or more milk reared under intensive management system with imported semen of progeny tested bulls), midterm (inseminate cross bred Holstein-Friesian cows yielding 6-10 kg milk reared under semi intensive management system with semen of progeny tested 50% Holstein-Friesian bulls) and long term goal (inseminate native cows reared under low input production system with semen of superior progeny tested/pedigree bulls of local

cattle for milk production also been set up by National Livestock Development Policy, Ministry of Fisheries and Livestock (NLDP, 2007). Considering the demerits of overall crossbreeding, selective breeding within the breeds is also recommended to conserve the cattle genetic resources. In both crossbreeding and selective breeding, animal selection is crucial for genetic improvement. However, identifying high-yielding animals remains a challenge, with conventional, time-consuming selection methods still in use. As a result, sustainable genetic improvement for high production has been limited over the past few decades. Recently, the Bangladesh Livestock Research Institute (BLRI) attempted to identify SNP markers for Bangladeshi cattle from genes previously reported in other cattle. This was the first effort toward marker-assisted selection for milk production traits in Bangladeshi cattle. However, they were unable to link the identified marker to phenotypic data.

Milk, a white opaque fluid rich in lipids, protein, lactose, and calcium, is often considered one of the best dietary sources of nutrition. It also contains a wide variety of micronutrients and other bioactive components produced by the mammary gland. These include vitamins, minerals, oligosaccharides, immunoglobulins, cytokines, antibodies, enzymes, enzyme inhibitors, growth factors, hormones, and antibacterial agents. Each component is crucial for the newborn's health and development, which is why milk and other dairy products are essential for a healthy, balanced diet (Wickramasinghe *et al.*, 2012; Pereira, 2014). The milk fat content and composition are key factors that determine the nutritional and technological quality of dairy products (Chilliard *et al.*, 2003). Fat content of milk is abundant in saturated fatty acids (SFA) and low in polyunsaturated fatty acids (PUFA). Therefore, fatty acid (FA) composition is valued economically, and enhancing milk FA composition, particularly by raising unsaturated fatty acid, is essential (Mannen, 2011). The fatty acid composition of milk is a heritable trait, with heritability ranging from 0.31 to 0.73. (Inoue *et al.*, 2008). Several recent studies have proposed genetic improvements to the nutritional quality of milk, focusing on its fatty acid profile (Abe *et al.*, 2009; Conte *et al.*, 2010; Matsumoto *et al.*, 2012; Mauric *et al.*, 2019).

To gain better insight into mammary biology and accelerate the rate of genetic gain in dairy cattle, numerous researchers have focused on identifying the genes and polymorphisms that influence bovine milk production. Fatty acid synthase, a multifunctional enzyme complex that controls the de novo synthesis of long-chain fatty acids, is a potential candidate gene for determining fat content in bovine milk and beef (Roy *et al.*, 2006; Schennink *et al.*, 2009; Matsumoto *et al.*, 2012; Li *et al.*, 2016; Mauriae *et al.*, 2017). The bovine *FASN* gene, spanning 19,770 bp and consisting of 42 exons and 41 introns, is located on BTA19 (Roy *et al.*, 2006; Kale *et al.*, 2021). It contains two main domains: the thioesterase (TE) and β -ketoacylreductase (KR) domains, which together produce long-chain fatty acids

of varying lengths, while the upstream Acyl carrier protein domain helps terminate chain elongation (Chakravarty *et al.*, 2004). The TE domain is thought to play a key role in regulating the fatty acid composition and content in bovine and other animals (Gibson *et al.*, 1958).

Till now, several studies have been conducted to explore the possible association between *FASN* gene polymorphisms and milk production traits. Following the identification of the *FASN* gene sequence and the structure of the fatty acid synthase complex, researchers have shifted their focus to investigating the association between polymorphisms *FASN* gene and their correlation with milk fatty acid composition (Kale *et al.*, 2021).

The *FASN* gene is found in a linkage region containing QTL for milk fat content and is a key enzyme for fatty acid synthesis, making it a promising candidate for milk production traits such as milk, protein, and fat yields, which are the primary breeding goals in dairy cattle selection. The present study aimed to screen SNPs in the *FASN* gene, analyze the protein's three-dimensional (3D) structure to understand the functional consequences of the polymorphism and evaluate its link to milk production traits in Bangladeshi local and Holstein cross cattle with the goal of potential application in cattle breeding.

Materials and Methods

The complete workflow used in this study is illustrated in Fig. 1




Fig. 1: The complete workflow used in this study

Ethical statement

All procedures involving animals and samples were approved (approval: NIBREC2017-02) by the Ethical Review Committee of the National Institute of Biotechnology (NIB), Bangladesh, where the experiment was conducted.

Experimental animals and phenotypic data collection

The National Institute of Biotechnology (NIB) ethical review committee in Bangladesh gave their clearance to the experiment (approval: NIBREC2017-02). A total of 100 crossbred F1 generation cows (Bangladeshi Local X Holstein) were selected from Central Cattle Breeding and Dairy Farm (CCBDF), Savar, Dhaka, where they were kept under the same management and environment condition. Milk samples were collected from each milking cow once during the whole lactation period, specifically between 90 and 100 days. The milk samples were then promptly sent to the Animal Biotechnology Division of the National Institute of Biotechnology (NIB) for phenotypic data generation (protein and fat percentage) using an auto milk analyzer (Lactoscan, Milk Analyzer, Bulgaria). Milk yield data was collected from CCBDF for each selected milking cow (305 days milk yield).

Screening polymorphisms and genotyping

DNA was extracted from collected blood samples (subset of cows, n=100) using the TIANamp Blood DNA Kit (TIANGEN BIOTECH (BEIJING) Co., Ltd.) according to the manufacturer's guidelines and instructions. Primers were designed and used based on Alim *et al.* (2013) and synthesized by Invitrogen (Invitrogen Life Technologies, China). A DNA pool was prepared with an equal volume and concentration of DNA samples taken from each animal (50 ng/ μ L/animal). Pool DNA was sequenced for preliminary screening of potential SNPs. After getting potential SNPs, a series of analyses was done to identify the most effective SNP. PCR amplifications were conducted with pooled samples using a programmable thermal cycler (Biometra GmbH, Germany). The 25 μ L reaction volume mixture contained 2 μ L DNA (50 ng), 1 μ L (1 μ M) each specific *FASN* gene forward and reverse primers, 8.5 μ L molecular grade water and 12.5 μ L of Invitrogen's DreamTaq Green PCR Master Mix. The amplification consisted of an initial denaturation at 94°C for 5 min, followed by 35 cycles of 94°C for 30 s, annealing at 58°C for 30 s, extension at 72°C for 30 s, and a final extension at 72°C for 7 min. Azure c150 gel imaging workstations were used for gel electrophoresis of PCR the products to confirm amplification. The gel was prepared by 2% agarose. After confirmation, the PCR products were sequenced using the ABI3500 sequencer (Applied Biosystems, USA) at the Molecular Biotechnology Division, NIB. Sequence chromatographs were then checked. If double peak is found in the same place of chromatographs, it is assumed to be polymorphism. After that, all sequence data was

analyzed to confirm the detected genetic polymorphisms using BioEdit Sequence Alignment Editor version 7.0.9.0 and ClustalW multiple sequence alignment programs. After structural and functional impact analysis of all identified SNPs, only polymorphism (g.17924 A>G) was genotyped using PCR and sequencing techniques in all selected animals.

Structural and functional impact prediction of *FASN* mutant protein

Identified non-synonymous polymorphisms in the *FASN* gene were examined for their possible effects on protein structure and function, which may affect the milk phenotype. To detect remote protein homology and predict structures, HHpred (<https://toolkit.tuebingen.mpg.de/tools/hhpred>) was used, while PROCHECK (<https://servicesn.mbi.ucla.edu/PROCHECK/>) was employed to assess structural integrity. The PDB-formatted three-dimensional model of the protein (Fig. 2) was created by the HHPred server. The PROCHECK server was used to produce a Ramachandran plot to verify the quality of the model. The Ramachandran plot and SAVES web-based server confirmed structure quality. The effect of SNPs on protein stability was predicted by MUpro (<https://www.ics.uci.edu/~baldig/mutation.html>) using Support Vector Machines and Neural Networks. Protein motion, stability and flexibility impact was analyzed by DynaMut2 (Rodrigues *et al.*, 2021). The ConSurf server was used for conservation profile analysis (Ashkenazy *et al.*, 2016), while domain identification was performed with the Pfam serve (Apweiler *et al.*, 2001). To check the impact of the mutations, the ConSurf web-based tool was employed to examine the evolutionary conservation of amino acid residues in the proteins. Molecular dynamics simulations of mutant *FASN* proteins and wild-type protein were performed using GROMACS for 100 ns (Abraham *et al.*, 2015).




Fig. 2: Three dimensional model of *FASN* mutant g.17924 G>A (T2264A). The position of the mutated amino acid is marked in red

Root Mean Square Deviation (RMSD) was calculated to assess structural changes in a molecule over time, particularly in proteins. Root Mean Square Fluctuation (RMSF) evaluated the regional flexibility of the protein, as mutations can either decrease or increase flexibility in previously stable regions. These flexible regions often play crucial roles in molecular recognition, binding, and enzymatic activity. A higher RMSF value indicates greater flexibility at a specific amino acid position. In molecular dynamics (MD) simulations, Solvent Accessible Surface Area (SASA) was employed to predict the impact of mutations on the stability of a protein's hydrophobic core. The radius of gyration was used to measure the protein's compactness, where a relatively stable value indicates proper folding, while fluctuations suggest unfolding. Mutations that increase the exposure of hydrophobic regions to the solvent (reflected by an increase in SASA) can destabilize the protein. SASA is a key property for understanding molecular interactions, stability, folding, and binding, as it reveals which regions of the molecule are solvent-exposed, buried, or involved in interactions.

Statistical analysis

MATLAB software (ver. 7.11.0.584) was used to trace pedigree data back one generation, a result total number of animals increased 300 for association analysis. POPGENE software (ver. 1.32) was used to perform the Hardy-Weinberg equilibrium test and to calculate allelic and genotypic frequencies at the loci. SAS software (ver. 9.1.0, SAS Institute Inc., USA) was used to estimate the effects of genotypes on milk production traits. The analysis was performed using the mixed procedure with an animal model (Lynch and Walsh, 1997):

$$Y = \mu + hys + L + G + \alpha + e$$

Where,

Y: Phenotypic value

μ : Average mean

hys: An effect of herd-year-season

L: Fixed lactation effect

G: Refer to fixed effect corresponding to the genotype of polymorphisms

α : Refer to random polygenic component for pedigree relationships

e: A random residual

The Bonferroni correction was applied to adjust the significance threshold for multiple comparisons, reducing the likelihood of false positives in multiple t-tests. This correction was calculated by dividing the significance level of a single test by the number of tests conducted. Since each trait had three genotype levels, three t-tests were performed. Consequently, the Bonferroni-adjusted significance thresholds were set at 0.0167 (0.05/3) and 0.0033 (0.01/3).

Least squares mean values were used for multiple comparisons to estimate the effects of *FASN* polymorphic genotypes on milk production traits, likely by comparing the mean values of different genotypic groups. Falconer and Mackay's equation (1996) was

applied to calculate the additive (a), dominance (d), and allele substitution (α_1) effects, providing insights into how specific alleles influence the phenotype. These effects were computed using the equations:

$$\text{Additive effect (a): } a = \frac{AA - BB}{2}$$

$$\text{Dominance effect (d): } d = AB - \left(\frac{AA + BB}{2} \right)$$

$$\text{Allele substitution effect } (\alpha_1): \alpha_1 = a + d(q - p)$$

Where,

AA and BB: Represent homozygous genotypes

AB: The heterozygous genotype

p and q: The allele frequencies

Results

Screening of single nucleotide polymorphisms and genotypes

Based on the bovine *FASN* sequence available in the GenBank database (accession No.: AF285607.2), the gene consists of 41 exons with a total length of approximately 19,760 base pairs. According to our sequencing result using previously reported primers, four SNPs were identified in this study: g.13965 C>T, g.16907 T>C, g.17924 A>G, and g.18663 T>C, located in exons 26, 36, 38, and 41, respectively. Among these, the g.17924 A>G polymorphism was predicted to alter the *FASN* protein by substituting threonine (ACC) with alanine (GCC). The remaining three SNPs (g.13965 C>T, g.16907 T>C, and g.18663 T>C) were silent mutations. The allelic and genotypic frequencies are summarized in Table 1. The Chi-square (χ^2) test confirmed that the genotypic frequencies of the loci were in Hardy-Weinberg equilibrium ($P > 0.05$) within the population, suggesting that selection pressure at these sites was not significantly strong.

Association analysis revealed that certain SNPs were strongly associated with three milk production traits, as indicated by raw P-values < 0.05 . However, these associations did not remain statistically significant after applying the Bonferroni correction for multiple t-tests (Table 2). Notably, the g.17924 A>G polymorphism exhibited a moderate association with milk yield. Based on these findings, a structural and functional evaluation of the mutant protein formed by the polymorphism g.17924 A>G. Cows carrying the G allele at this locus showed dominance in milk yield within the population, potentially increasing production by 338.91 kg over a full lactation period (Table 2).

Structural and functional impact prediction of *FASN* mutant protein associated with milk traits

Structural impact prediction

The results from HHPred and the PROCHECK server indicated that most residues in the three-dimensional models, for both the wild-type and mutant forms, were

located in the most favored regions (Fig. 3).

Conservation profile analysis

The ConSurf web tool results were displayed as a structural representation of the protein sequence, highlighting the predicted structural and functional

residues. In this analysis, the 2264th position in FASN showed an average conservation profile (Fig. 4).

Identification of domain

The FASN mutant T2264A was located within the thioesterase (TE) domain of the FASN protein (Fig. 5).

Table 1: Genotypic and allelic frequencies and Hardy-Weinberg equilibrium χ^2 test of FASN genotypes

Polymorphisms	Genotypic frequency			Allelic frequency		Hardy-Weinberg equilibrium χ^2 test
g.17924 G>A	GG	AG	AA	G	A	P>0.05
	0.61	0.34	0.05	0.75	0.25	

Table 2: Least squares mean (LSM) and standard errors (SE) for milk production traits of different *FASN* genotypes and Additive and allele substitution effects of SNPs on milk production traits in Bangladeshi local and Holstein cross cattle

Locus	Genotype	Milk yield (kg)	Fat percentage (%)	Protein percentage (%)	Additive (a), dominant (d) and allele substitution (α 1) effects		
					Milk yield (kg)	Fat percentage (%)	Protein percentage (%)
g.17924 G>A	AA	2156.19±268.428	3.67±0.169	3.11±0.058	261.295 (a)* G>A	0.044 (a)	-0.007 (a)
	AG	1754.45±124.258	3.67±0.078	3.07±0.027	-140.447 (d)	0.041 (d)	-0.043 (d)
	GG	1633.60±94.903	3.59±0.059	3.12±0.020	338.906 (α 1)*	0.021 (α 1)	0.016 (α 1)
P-value		<0.0001 (0.22-0.54 corrected)	<0.0001 (0.90 corrected)	<0.0001 (0.90 corrected)			

* Significant additive effect or allele substitution effect at P<0.05 level




Fig. 3: Ramachandran plot of FASN mutant g.17924 G>A (T2264A) generated by PROCHECK

The TE domain is a structural and functional motif commonly found in enzymes involved in lipid metabolism, biosynthesis, and degradation.

Interatomic interactions prediction

MUpro analysis indicated that the T2264A mutation in FASN reduced protein stability, with a ΔG value of -1.4982588 kcal/mol. Similarly, DynaMut2 predicted a stability change ($\Delta\Delta G_{\text{Stability}}$) of -0.13 kcal/mol for the same mutation. A negative $\Delta\Delta G$ value suggests that the mutation decreases protein stability (Fig. 6).

Molecular dynamic simulation

Variations in RMSD values indicate conformational changes in the protein. A significant increase in RMSD suggests major structural alterations, such as domain movements, loop motions, or unfolding events. The RMSD values of the wild-type and the T2264A mutant began to diverge at the 15 ns mark of the simulation. From that point onward, the T2264A mutant consistently

exhibited a lower RMSD profile compared with the wild type (Fig. 7).

In our study, RMSF analysis revealed that in case of the wild FASN and the T2264A mutant different regions were more flexible. The region around the 25th and the 120th residues were significantly more flexible in the mutant compared with the wild type whereas the regions close to 60th, 90th, 210th, and 250th residues were more flexible in the wild type (Fig. 8). In molecular dynamics, RMSD gives insight into overall structural stability, while RMSF reveals local flexibility and dynamic behavior. In our study, we calculate RMSD and RMSF for better understanding of FASN T2264A mutant protein. Radius of gyration analysis revealed structural differences between the wild-type and mutant FASN (Fig. 9). A higher SASA value indicates an increased likelihood of protein destabilization due to greater solvent accessibility. Although the T2264A mutant initially maintained a lower SASA value up to 20 ns, it remained higher than the wild type thereafter, suggesting




Fig. 4: Conservation profile analysis of FASN protein

a higher probability to solvent-induced disruption (Fig. 10).

Discussion

Since 1961, CCBS has conducted crossbreeding experiments with cattle across eight pure breeds:

Bangladeshi Local, Sahiwal Kenyan, Sahiwal, Red Sindhi, Friesian, Jersey, Holstein-Friesian and Tharparker, along with their crosses. After the establishment of CCBS, cow information has been recorded consistently. The earmark numbers (1 to 9,999) of the first batch was completed by August 26, 1983, followed by a second batch starting the next day. All




Fig. 5: FASN Domains




Fig. 6: Interatomic interactions of FASN wild-type and T2264A mutant revealed by dyanmute2 analysis




Fig. 7: RMSD of the wild-type (green) FASN and the T2264A mutant (red). The X-axis represents the time (ns) while the Y-axis represents the RMSD value (nm)




Fig. 8: RMSF of the wild-type (green) FASN and the T2264A mutant (red). The X-axis represents the amino acid residues while the Y-axis represents the RMSF value (nm)




Fig. 9: Radius of gyration of the wild-type FASN (green) and the T2264A mutant (red). The X-axis represents the time (ps) while the Y-axis represents the Area (nm²)




Fig. 10: SASA of the wild-type FASN (green) and the T2264A mutant (red). The X-axis represents the time (ps) while the Y-axis represents the SASA value (nm)

animals at CCBS are fed and managed under standard conditions throughout the year. Concentrate feeds, including rice polish, khesheri, wheat bran, till oil cake, and salt, are provided twice daily, before milking started at morning and in the evening, based on body

requirements. Green grasses, like maize, napier, para, and oats, are supplied year-round. Cows are milked twice daily, machine milking for high-yielding cows and with hand-milking for low-yielding ones (Hossain *et al.*, 2002). In our study, two conditions have been included:

- a) Crossbred F1 generation (Bangladeshi Local X Holstein) cows
- b) Reared in the same management situation

Although CCBS has a total of 14,512 records for all animals (Hossain *et al.*, 2002; Hamid *et al.*, 2017) but after setting the condition sample size become lower.

Since the 1930s, various scattered efforts have been made to improve cattle production by introducing foreign genes, but with no notable success. Key challenges include low-quality breeds, disease outbreaks, insufficient vaccines, feed scarcity and high costs and fluctuating market prices. Science-driven cattle breeding in Bangladesh remains underdeveloped, lacking a defined national strategy or vision (Hamid *et al.*, 2017). Molecular genetics techniques are commonly applied to pinpoint genes associated with economically advantageous traits in dairy cattle. These genes are then used as selection markers in marker-assisted selection (MAS) breeding, thus driving advancements in the dairy cattle industry. In essence, breeding programs aim to pinpoint superior genotypes for economically valuable traits by leveraging data on animal performance, familial relationships, and molecular information. This enables the propagation of favorable genes throughout the population (Yudin and Voevoda, 2015; Miglior *et al.*, 2017). In the present study, *FASN* gene had been chosen as positional candidate aiming to find out a suitable marker that could be used in cattle breeding. We identified mutation in *FASN* (SNP g.17924 A>G) that showed an association with milk yield trait in our population.

In this study, the frequency of A allele (0.25) at g.17924 A>G, in our population was higher than that found by (Bhuiyan *et al.*, 2009) and (Maharani *et al.*, 2012) in the Hanwoo population (0.16 and 0.19, respectively), but lower than those reported in Angus beef cattle (0.62) (Zhang *et al.*, 2008), 0.31 in Friesian cattle (Morris *et al.*, 2007) and 0.53 in Dutch Holstein-Friesian population (Schennink *et al.*, 2009) 0.54 in Canadian Angus and Charolais-based commercial crossbred steers (Li *et al.*, 2011). Very recent study on Polish Red (RP), Polish Red-and-White (ZR) and Polish Holstein-Friesian Red-and-White (RW) breeds, SNP g.17924 A/G, the GG genotype was most frequent in ZR and RP cows, with frequencies of 0.73 and 0.82, respectively ($P < 0.01$) and RW cow had the highest frequency (0.30) of the A/G genotype ($P < 0.01$) (Przybylska and Kuczaj, 2024). These discrepancies might be induced by long term cross breeding for milk yield, natural selection, and random drift. Many previous studies with QTL mapping and candidate gene approach found significant associations of *FASN* gene with milk fatty acid composition (Morris *et al.*, 2007; Schennink *et al.*, 2009; Stoop *et al.*, 2009) and milk fat traits (Roy *et al.*, 2006; Ordovas *et al.*, 2008; Schennink *et al.*, 2009)

in different cattle populations, but in our population the significant single locus associations showed a clear effect additive and allele substitution effects on milk yield trait, whereas it did not reach significance for other milk traits. The inconsistency may result from interactions with background genes in various cattle breeds. In general, the effects of polymorphism can vary across populations or breeds due to their unique genetic backgrounds. Bangladeshi Holstein cross cattle have developed through crossbreeding over the past 70 years, involving non-descriptive Deshi cattle and purebred Holstein bull semen from Australia, America, or Europe (NLDP, 2007). Morris *et al.* (2007) identified five SNPs in the *FASN* gene including g.17924 A>G, a non-synonymous SNP leading to a p.Tyr>Ala amino acid change. Our findings were consistent with the current literature on amino acid replacement.

The mutational impact analysis revealed that the mutation in the *FASN* (SNP g.17924 A>G) gene likely destabilizes the protein, affecting its functionality. The function, regulation and activity of a given protein depend on how well its structure holds together. Proteins degrade, misfold, and clump when stability decreases, eventually becoming dysfunctional (Rozario *et al.*, 2021). Analysis suggests the *FASN* mutant T2264A changes in interatomic interactions and protein stability. The *FASN* mutant T2264A was in the conserved thioesterase domain. The thioesterase domain is a conserved structural motif present in enzymes known as thioesterases. These enzymes essential for catalyzing the hydrolysis of thioesters, which are chemical compounds with a sulfur atom bonded to an acyl group that are involved in various biological processes, including fatty acid metabolism, polyketide biosynthesis, and protein modification. Molecular dynamics simulations show clear differences between the mutant and wild type *FASN* in terms of RMSF, RMSD, Radius of gyration, and SASA profile. In summary, the functional and structural analysis of SNP g.17924 A>G mutant protein showed clear differences from wild type protein and may have significant implications for protein structure and function, which could affect various biological processes.

Furthermore, our findings indicate that SNP in the bovine *FASN* gene is linked to variations in milk production traits, reinforcing previous evidence that *FASN* may be the causative gene for QTL associated with fat-related traits in dairy cattle. Structural and functional analysis of the mutation in *FASN* (SNP g.17924 A>G) also supports the association. However, additional experimental validation is required to confirm these findings.

The present study revealed that nsSNPs (SNP g.17924 A>G) was significantly associated with milk yield in the studied population. Such significant effect was possible due to amino acid replacement in *FASN* protein for this nsSNPs that might change the phenotypes. In comparison with G allele at g.17924 A>G position, an allele could increase the milk production 338.9 kg in a full lactation period. Structural prediction,

domain identification, conservation profile, interatomic interactions and molecular dynamic simulation analysis also support the result. These results, along with the functional and structural evaluation of both the *FASN* wild-type and mutant proteins, provide valuable insights that, with proper validation, could be applied in selective breeding programs to enhance milk production performance in dairy cattle in Bangladesh.

Acknowledgements

The authors would like to acknowledge the Department of Livestock Services, Bangladesh for supporting this study. The study was funded by the Bangladesh Academy of Sciences under BAS-USDA program Code No. LS-15/2017. The funder has no role in the design and conduct of the study; collection, management, analysis and interpretation of the data, preparation, review and approval of the manuscript; or decision to submit the manuscript for publication. The authors also would like to thank a post-doctoral researcher (native speaker), Wuhan University, for his assistance with English expression and editing.

Conflict of interest

The authors declare no conflicts of interest.

References

- Abe, T; Saburi, J; Hasebe, H; Nakagawa, T; Misumi, S; Nade, T; Nakajima, H; Shoji, N; Kobayashi, M and Kobayashi, E (2009). Novel mutations of the *FASN* gene and their effect on fatty acid composition in Japanese Black beef. *Biochem. Gen.*, 47: 397-411.
- Abraham, MJ; Murtola, T; Schulz, R; Pall, SR; Smith, JC; Hess, B and Lindahl, E (2015). GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX*. 1: 19-25.
- Alim, MA; Wang, P; Wu, XP; Li, C; Cui, XG; Zhang, SL; Zhang, Q; Zhang, Y and Sun, DX (2013). Effect of *FASN* gene on milk yield and milk composition in the Chinese Holstein dairy population. *Anim. Gen.*, 45: 111-113.
- Apweiler, R; Attwood, TK; Bairoch, A; Bateman, A; Birney, E; Biswas, M; Bucher, P; Cerutti, L; Corpet, F and Croning, MDR (2001). The InterPro database, an integrated documentation resource for protein families, domains and functional sites. *Nucl. Aci. Res.*, 29: 37-40.
- Ashkenazy, H; Abadi, S; Martz, E; Chay, O; Mayrose, I; Pupko, T and Ben-Tal, N (2016). ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation in macromolecules. *Nucl. Aci. Res.*, 44: W344-W350.
- Bhuiyan, MSA; Yu, SL; Jeon, JT; Yoon, D; Cho, YM; Park, EW; Kim, NK; Kim, KS and Lee, JH (2009). DNA polymorphisms in *SREBF1* and *FASN* genes affect fatty acid composition in Korean cattle (Hanwoo). *Asi-Aus. J. Anim. Sci.*, 22: 765-773.
- Chakravarty, B; Gu, Z; Chirala, SS; Wakil, SJ and Quijcho, FA (2004). Human fatty acid synthase: structure and substrate selectivity of the thioesterase domain. *Proc.*

- Nat. Acad. Sci., 101: 15567-15572.
- Chilliard, Y; Ferlay, A; Rouel, J and Lamberet, G** (2003). A review of nutritional and physiological factors affecting goat milk lipid synthesis and lipolysis. *J. Dairy Sci.*, 86: 1751-1770.
- Conte, G; Mele, M; Chessa, S; Castiglioni, B; Serra, A; Pagnacco, G and Secchiari, P** (2010). Diacylglycerol acyltransferase 1, stearoyl-CoA desaturase 1, and sterol regulatory element binding protein 1 gene polymorphisms and milk fatty acid composition in Italian Brown cattle. *J. Dairy Sci.*, 93: 753-763.
- DLS and Director, G** (2021). <https://www.dls.gov.bd/site/page/22b1143b-9323-44f8-bfd8-647087828c9b/Livestock-Economy>. PP: 1-5.
- Falconer, DS and Mackay, TFC** (1996). *Introduction to quantitative genetics*. 4th Edn., New York, Longman Scientific and Technical. PP: 24-47.
- Gibson, DM; Titchener, EB and Wakil, SJ** (1958). Studies on the mechanism of fatty acid synthesis V. Bicarbonate requirement for the synthesis of long-chain fatty acids. *Biochim. Biophys. Act.*, 30: 376-383.
- Hamid, M; Rahman, A; Zaman, MA and Hossain, KM** (2017). Cattle genetic resources and their conservation in Bangladesh. *Asian J. Anim. Sci.*, 11: 54-64.
- Hossain, KB; Takayanagi, S; Miyake, T; Moriya, K; Bhuiyan, A and Sasaki, Y** (2002). Statistical genetic studies on cattle breeding for dairy productivity in Bangladesh: I. Genetic improvement for milk performance of local cattle populations. *Asi-Aus. J. Anim. Sci.*, 15: 627-632.
- Inoue, K; Shoji, N and Kobayashi, M** (2008). Genetic relations among fat melting point, fatty acid composition and carcass traits in Japanese Black cattle. *Anim. Sci. J. (Japan)*, 79: 1-8.
- Kale, DS; Singh, J; Sathe, YB and Patil, DV** (2021). *FASN* gene and its role in bovine milk production. *Int. J. Biotech. Trends Technol.*, 11: 20-25.
- Li, C; Aldai, N; Vinsky, M; Dugan, MER and McAllister, TA** (2011). Association analyses of single nucleotide polymorphisms in bovine *stearoyl-CoA* desaturase and fatty acid synthase genes with fatty acid composition in commercial crossbred beef steers. *Anim. Gen.*, 43: 93-97.
- Li, C; Sun, D; Zhang, S; Yang, S; Alim, MA; Zhang, Q; Li, Y and Liu, L** (2016). Genetic effects of *FASN*, *PPARGC1A*, *ABCG2* and *IGF1* revealing the association with milk fatty acids in a Chinese Holstein cattle population based on a post genome-wide association study. *BMC Genet.*, 17: 1-16.
- Lynch, M and Walsh, B** (1997). *Genetics and analysis of quantitative traits*. 1st Edn., Sunderland, MA, USA, Sinauer Associates, Inc., PP: 35-48.
- Maharani, D; Jung, Y; Jung, WY; Jo, C; Ryoo, SH; Lee, SH; Yeon, SH and Lee, JH** (2012). Association of five candidate genes with fatty acid composition in Korean cattle. *Mol. Bio. Rep.*, 1: 1-9.
- Mannen, H** (2011). Identification and utilization of genes associated with beef qualities. *Anim. Sci. J.*, 82: 1-7.
- Matsumoto, H; Inada, S; Kobayashi, E; Abe, T; Hasebe, H; Sasazaki, S; Oyama, K and Mannen, H** (2012). Identification of SNPs in the *FASN* gene and their effect on fatty acid milk composition in Holstein cattle. *Liv. Sci.*, 144: 281-284.
- Mauriaie, M; Masek, T; Beniae, M; Spehar, M and Stareeviae, K** (2017). Effect of *DGAT1*, *FASN* and *PRL* genes on milk production and milk composition traits in Simmental and crossbred Holstein cattle. *Ind. J. Anim. Sci.*, 87: 859-863.
- Mauric, M; Masek, T; Ljoljic, DB; Grbavac, J and Starcevic, K** (2019). Effects of different variants of the *FASN* gene on production performance and milk fatty acid composition in Holstein Simmental dairy cows. *Vet. Med.*, 64: 101-108.
- Miglior, F; Fleming, A; Malchiodi, F; Brito, LF; Martin, P and Baes, CF** (2017). A 100-year review: Identification and genetic selection of economically important traits in dairy cattle. *J. Dairy Sci.*, 100: 10251-10271.
- Morris, CA; Cullen, NG; Glass, BC; Hyndman, DL; Manley, TR; Hickey, SM; McEwan, JC; Pitchford, WS; Bottema, CDK and Lee, MAH** (2007). Fatty acid synthase effects on bovine adipose fat and milk fat. *Mamm. Geno.*, 18: 64-74.
- NLDP and Director, G** (2007). National livestock development policy, Ministry of Fisheries and Livestock, Government of the People's Republic of Bangladesh. PP: 1-115.
- Ordovas, L; Roy, R; Pampin, S; Zaragoza, P; Osta, R; Rodriguez-Rey, JC and Rodellar, C** (2008). The g.763 G>C SNP of the bovine *FASN* gene affects its promoter activity via Sp-mediated regulation: implications for the bovine lactating mammary gland. *Physio. Geno.*, 34: 144-148.
- Pereira, PC** (2014). Milk nutritional composition and its role in human health. *Nutrition*. 30: 619-627.
- Przybylska, P and Kuczaj, M** (2024). The effects of two selected single nucleotide polymorphisms of the fatty acid synthase gene on the fat content and fatty acid profile of cow's milk from the Polish Holstein Friesian Red-and-White breed versus two Polish Red-and-White and Polish Red conservation breeds kept in Poland. *Animals*. 14: 2268-2275.
- Rodrigues, CHM; Pires, DEV and Ascher, DB** (2021). DynaMut2: Assessing changes in stability and flexibility upon single and multiple point missense mutations. *Prot. Sci.*, 30: 60-69.
- Roy, R; Ordovas, L; Zaragoza, P; Romero, A; Moreno, C; Altarriba, J and Rodellar, C** (2006). Association of polymorphisms in the bovine *FASN* gene with milk-fat content. *Anim. Gen.*, 37: 215-218.
- Rozario, LT; Sharker, T and Nila, TA** (2021). *In silico* analysis of deleterious SNPs of human *MTUS1* gene and their impacts on subsequent protein structure and function. *PloS One*. 16: e0252932-e0252945.
- Schennink, A; Bovenhuis, H; Leon Kloosterziel, KM; Van Arendonk, J.A.M and Visker, MHPW** (2009). Effect of polymorphisms in the *FASN*, *OLR1*, *PPARGC1A*, *PRL* and *STAT5A* genes on bovine milk fat composition. *Anim. Gen.*, 40: 909-916.
- Stoop, WM; Schennink, A; Visker, M; Mullaart, E; Van Arendonk, JAM and Bovenhuis, H** (2009). Genome-wide scan for bovine milk-fat composition. I. Quantitative trait loci for short- and medium-chain fatty acids. *J. Dairy Sci.*, 92: 4664-4675.
- Wickramasinghe, S; Rincon, G; Islas-Trejo, A and Medrano, JF** (2012). Transcriptional profiling of bovine milk using RNA sequencing. *BMC Geno.*, 13: 1-14.
- Yudin, NS and Voevoda, MI** (2015). Molecular genetic markers of economically important traits in dairy cattle. *Rus. J. Gen.*, 51: 506-517.
- Zhang, S; Knight, TJ; Reecy, JM and Beitz, DC** (2008). DNA polymorphisms in bovine fatty acid synthase are associated with beef fatty acid composition. *Anim. Gen.*, 39: 62-70.